

National Journal of Research in Ayurved Science

# Randomised single blind pilot clinical study of *kharjura* on haemoglobin percentage with special reference to *sushruta's rakta dhatu poshankal.*"

Supriya P Tonge<sup>\*1</sup>, Swati Chobhe<sup>2</sup>, Manisha B Walunj<sup>3</sup>

- 1. Assistant Professor, KriyaSharira, SST Ayurved college, Ghulewadi, Sangamner, Maharashtra, 422605
- 2. Associate Professor, Kriya Sharira, P.D.E.A S Ayurved College and Research Centre, Nigdi, Pune, Maharashtra, Email: <a href="mailto:swatichobhe@yahoo.com">swatichobhe@yahoo.com</a>; 9850716124
- Assistant Professor, Rasashastra and Bhaishajya kalpana, SST Ayurved college, Ghulewadi ,Sangamner, Maharashtra, 422605 Email: drmanishawalunj@gmail.com; 8460150540

\*Corresponding author: Email: <a href="mailto:suprivatonge1@gmail.com">suprivatonge1@gmail.com</a> ; 8308252487

# **ABSTRACT:**

Introduction: Amongst Sapta Dhatu, Rakta Dhatu is of prime importance as it is mainly responsible for Jeevan Karma and Prananuvartana. According to Dhatu Poshana Kala theory by Acharya Susruta, *Rakta Dhatu* is formed on the 5<sup>th</sup> day after formation of AharaRasa. Rakta Kshaya can be correlated with symptoms of anaemia due to malnutrition. Kharjura (Phoenix dactylifera, Linn.) now days well known as dates is of neutraceutical value has been used in treatment of Rakta Dhatu since ages. Aim: To assess effect Of *Khajura* with Dhatriloha and Dhatriloha alone on haemoglobin percentage with special

reference to Sushruta's Rakta Dhatu J-R A S Poshankal."Present pilot study was performed to set guidelines with special reference to treatment and analysis period for chief clinical trial and taking into consideration Dhatu Poshana Kala Theory screening of patients for Hb % was done at intervals of 5 days. Materials and methods: Kharjura Laddu was prepared, patients fulfilling inclusion criteria of Rakta Kshaya were administered with Kharjura *Laddu*(10g)+*Dhatri Loha*(250mg) and Dhatri Loha(250mg) alone in groups A and B respectively for 10 days in Rasayan *Kala*(early in morning).**Results:** Kharjura Laddu+Dhatri Loha group revealed better results in relieving symptoms and increasing Hb%. Both groups exhibited increase in Hb% at 5 days interval after start of treatment. **Conclusion:** *Kharjura* can be used as neutraceutical along with *Dhatri Loha* in management of *Rakta Kshaya*. Insignificant increase in Hb% at interval of 5 days of consumption of drug ascertains *Dhatu Poshan Kala* theory.

**Keywords:** *Kharjura*, neutraceutical, Hb%, *Dhatuposhana Kala* theory.

## **INTRODUCTION:**

*Dosha, Dhatu & Mala* are considered as three basic elements of body. The conceptof *Sapta Dhatu* of *Ayurveda* refers to the physiology of basic nutritional and structural factors of the body. They are being constantly formed, destroyed and reformed with appropriate material derived from *Poshaka Dravya* (nutrient substances) present in food from the time of conception to death.

Dosha controls all physiological activity while Dhatu perform Dharana *Karma*(sustaining action). There are seven Dhatu viz. Rasa, Rakta, Manmsa, Meda, Majja and Shukra. Out of these seven Dhatu, Rakta Dhatu has prime importance because it is mainly responsible for Jeevan Karma and Prananuvartana<sup>1</sup>. According to modern science, haemoglobin is the protein in

erythrocytes that is responsible for the delivery of oxygen from the lungs to the tissues, and the transport of carbon dioxide from the tissues back to the lungs.This process is similar with **Prananuvartan** function of *Rakta Dhatu.*<sup>2</sup> All ancient Samhita have postulated the view that Sapta Dhatu are produced in a progressive metamorphosis, kind of beginning with the Rasa Dhatu and ending with Shukra Dhatu. The previous Dhatu being transformed in to the next higher this theory knows as Dhatu -Poshankal. According to Acharva Susruta, by this rule Rakta Dhatu is formed on the 5<sup>th</sup> day after formation of AharaRasa.<sup>3</sup>

Biosynthesis of Rakta Dhatu depends on nutritional supplements from food and if these supplements are not received then subjected Rakta person get to Kshaya(deficiency). Rakta Kshaya can be correlated with symptoms of anaemia due to malnutrition. It is the most common nutritional deficiency disorder in the world. World Health Organisation has defined anaemia as a condition in which the haemoglobin content of blood is less than normal, due to deficiency of one or more essential nutrients, regardless cause of such deficiency. A haemoglobin level of 10 to 11 g/dl has been defined as early anaemia<sup>4</sup> Anaemia is established if the haemoglobin is below the cut-off points

recommended by W.H.O.<sup>5</sup> in such condition it is important for people to increase their consumption of iron-rich foods in diet such as dates (which is cheaper), red meat, fish, poultry and legume. *Kharjura (Phoenix dactylifera,* Linn.) now days well known as dates has been used in treatment of *Rakta Dhatu* since ages.

According to Ayurveda all the drugs are Panchabhautic and thus no Dravya are without medicinal property. Logically and intelligently every Dravya in this world can be used as a food and medicine, the same concept is coming forward in medicine which is called as nutraceuticals<sup>6</sup>.Now days dates can be called as a nutraceuticals.<sup>7</sup>*Kharjura* is the wonderful package of deliciousness and nutrition liked by everyone.

Primary aim of this particular trial is "to assess effect Of *Khajura* with *Dhatriloha* and *Dhatriloha* alone on haemoglobin percentage with special reference to *Sushruta's Rakta Dhatu Poshankal.*"

Present pilot study was performed to set guidelines with special reference to treatment and analysis period for chief clinical trial .Pilot study was carried out to assess effect of *Kharjura* intake on Hb% in management of anaemia and taking into consideration *Dhatu Poshana Kala Theory* screening of patients for hb % was done at intervals of 5 days to see whether there is increase in hb% after five days.

## AIM AND OBJECTIVES

To compare the efficacy of *Kharjura* as a neutraceutical with *Dhatriloha* to increase Haemoglobin Percentage.

## MATERIALS AND METHODS

# Materials -

- Kharjura Laddu Self prepared by investigator.
- Haemoglobin percentage Sahli's Hemoglobinometer.

# Material details of both drugs:-JRAS a) Kharjura –

Latin Name - Phoenix dactylifera, Linn.

#### **Classification**

- Kingdom Plantae
- Subkingdom Tracheobionta
- Superdivision Spermatophyta
- Division Magnoliophyta
- Class Liliopsida
- Subclass Arecidae
- Order Arecales
- Family Arecaceae
- Genus Phoenix
- Species dactylifer

## Nutritional values - As per USDA

(United States Department of

Agriculture), nutritional values of date fruit- *Kharjura (Phoenix dactylifera* Linn.) per 100 gm. are as follows<sup>8</sup>

Method of preparation of *Kharjura Laddu* :-

Preparation – Self prepared by investigator.

Ingredients: *Kharjura* (marketed by lion company ISO certified and FSSI certified)

Procedure -

- a) Manually removal of seed from *Kharjura*.
- b) Grinding of *Kharjura* in mixer grinder.
- c) Obtaining *Kharjura* paste and then lastly prepared *Kharjura* laddu (10 g)
- b) *Dhatriloha*<sup>9</sup>

Market\_sample of *Dhatri Loha* (tab-250 mg) manufactured by Baidyanath was used for present study.

# **Selection of patients:**

The study was conducted at SST Ayurved College, Sangamner after obtaining approval from the Institutional Ethics Committee (ks-1/33/Ethics/2017- 2018 dated-14/02/2018).

**Inclusion criteria:** Female students of 20-24yr age group with haemoglobin

level < 11 gm and >10 gm with prior informed consent were included in study.

#### **Exclusion criteria:**

While pregnant and lactating women, *Rakta-Kşhaya* due to the bleeding disorders, diagnosed with haematological disorders or anaemia due to aplastic anaemia, pernicious anaemia, haemolytic anaemia and other haemoglobinopathies were excluded from study.

#### Posology

The study was randomized, single blind study. Lottery method was adopted for random allocation of patients and the ones satisfying the inclusion criteria were grouped into two groups, namely, A and B.Patients in group A were administered with *Kharjura Laddu* (10 g) with Dhatriloha (250mg tab) while patient in B group were administered only Dhatriloha (250mg tab) in Prabhat Kala (early in morning) for period of 10 days. Patients were allowed to follow normal routine diet and regimen during treatment period. After completion of study, data were collected and analyzed statistically. Laboratory investigations

Haemoglobin percentage by Sahli's Hemoglobinometer<sup>10</sup> method was carried out at the time of screening of patients for inclusion and at the interval of 5 days and at 10 days(completion of treatment

period) respectively to assess effect of both the test drugs.

#### Assessment criteria

The assessment of patients was done before starting treatment, at interval of 5 days and after completion of treatment (10 days). Symptoms of Rakta Dhatu Kshaya described by Acharya Sushruta-Amla-sheeta Pratharna (craving for sour and cold substances), Twaka Parushya (dryness of skin) and Sira Shaithilya (loss of tone of veins) were considered as subjective parameters. Gradations of the symptoms were done by giving them score. This Scoring pattern [Table no.3] was used to assess the response of trial drugs objectively and for statistical analysis. Haemoglobin percentage was considered as objective criteria and analysed before treatment, at 5 days and after completion of the treatment (10 days).

#### **Data analysis**

The data obtained in clinical study was subjected to statistical tests and calculated on the basis of percentage of improvement in each parameter in all the treated groups. Statistical tests were performed by using GraphPad Prism 6 Software (Graphapad software Inc, La Jolla, CA 92037 USA) and sigma stat <sup>11</sup> Wilcoxson signed rank test was used to evaluate effect of test drugs in subjective parameters (p<0.05) in individual group. Effect on Hb % by both the drugs is assessed by paired t test (p<0.05). Comparison of relief in symptoms and improvement in Hb% by test drugs is done by applying CV (coefficient of variation) test to statistical data. Overall effect of test drugs was compared by unpaired t test (p<0.05).

#### **RESULTS:**

In total ten patients were registered (five in each group) for the pilot study. Five patients registered in each group completed treatment. At the end of treatment period data were collected and analysed stastically. (dia1: consort chart)

After completion of treatment period both test drugs produced insignificant relief (p>0.05) in symptoms – *Twaka Parushya* (dryness of skin) and *Amla Sheeta Pratharna* (craving for sour and cold substances). However percentage of relief was observed to be more in group B.(Table no:4 and table no.5)

In both treated groups increase in Hb % was noted at five days as well as ten days interval. Percentage increase in Hb % parameter was observed to be more in group B which was statistically insignificant (p>0.05). (Table no.6 and table no.7)

On comparing effect of drugs by CV(coefficient of variation) test it is observed that relief obtained in *Amla Sheeta Pratharna*(craving for sour and cold substances) was equal in both groups. Group B produced better relief in *Twaka Parushya* (dryness of skin) symptom and improvement in hb%. (table no.8)

When overall effect of test drug was compared by unpaired t test it was statistically insignificant (p>0.05)(table no.9).

# **DISCUSSION:**

Above results implies that, at the end of treatment period both the test drugs were effective in producing relief in symptoms of Rakta Kshaya described by Acharya Sushruta- Twaka Parushya (dryness of skin) and Amla Sheeta Pratharna (craving for sour and cold substances). At statistical level improvement was insignificant in both groups (p>0.05). However percentage relief was noted maximum in group B where patients were treated with Dhatri Loha and Kharjura Laddu in comparison to group A in which patients were administered with Dhatri Loha alone. CV evaluation also demonstrated group В better in

management of symptoms of Rakta Kshaya.

To assess effect of Kharjuara on Hb% improvement, Hb% was carried out at 5 days in accordance with Dhatu Poshana Kala theory and at the completion of treatment period (10 days). In both the treated groups apparent insignificant (p>0.05) increase in Hb% was observed at both five days and ten days interval. Result in Hb% signifies that both test drugs have additive effect on Hb%. As increase in Hb% in both treated groups is noted at five days interval it asserts Dhatu Poshankala theory that after consumption of drug process of formation of Rakta Dhatu is initiated and is evident within five days. Results of CV evaluation and percentage relief indicates group B better in improvement in hb% parameter where patients have been treated with Dhatri Loha along with Kharjura Laddu.

Overall results implies group B better in management of *Rakta Kshaya* and improvement in Hb% thus elucidating neutraceutical value of *Kharjura* with *Dhatri Loha* in increasing Hb%. However to form concrete conclusive remark study population should be large enough which will be done in main clinical study.

#### Probable mode of action of drug:

In context of avurveda science to increase Rakta Dhatu, one has to consume food and nutrients which has properties homologous to Rakta Dhatu. Rakta Dhatu and Kharjura both pursue similar properties (Guna Samanya). Both Rakta Dhatu and Kharjura have properties like Guru(heavy)and *Snigdha*(unctuous), Madhura(sweet) Rasa and it has got Vipaka. Kharjura Madhura acts as  $RaktaPittaprasadak^{12}$  hence it helps in biosynthesis of Rakta Dhatu quantitatively. Nutritional screening shows that the Kharjura contains 34mg/100gm of calcium. 6 g/100gm of iron and 30mg/100gm of vitamin C which are essential for formation of haemoglobin.<sup>13</sup> According to review of the phytochemical constituents of Kharjura mentioned by the phytochemists of 20th century, it contains all the necessary materials such as iron, vitamin C, etc. which are required erythrocyte<sup>14</sup>. the formation of for Kharjura is having erythrogenic properties i.e., it enhances production of red cells; facilitates the absorption of iron in the body there by increases the haemoglobin level.<sup>15</sup>

#### CONCLUSION

*Kharjura* can be used as neutraceutical along with *Dhatri Loha* in management of Rakta *Kshaya*. Pilot study revealed

insignificant increase in Hb% at interval of 5 days of consumption of drug ascertaining *Dhatu Poshan Kala* theory. ACKNOWLEDEMENT

SST Ayurveda college, Sangamner.

P.D.E.A.S Ayurved college and research centre, Nigadi Pune.

#### **REFERENCES:**

- Dr. Brahmanand Tripathi and Dr. Ganga sahay pandey,editors. Charaka Samhita of Agnivesha, Sutrasthan. Chapter 24. Reprint ed. Varanasi (India): Chaukambha Orientalia Publication; 2009. p. 124
- Dr. Ghanekar, Commentator. Susrutha Samhita, Sutrasthan. Chapter 14. Delhi(India): Maeherchandlachmandas
- Publication ; 2010. p. 69

2010.p.377

- Acharya JT, Editior.Susrutha Samhita,Sutrasthan.Chapter 14.Varanasi(India):Chaukambha Sanskrit Sansthan Publication;
- Mohan Harsh. Editior.Text book of pathology.Chapter 8 .New Delhi: Jaypee Brothers Medical Publishers ltd; 2010. p. 288-9
- Dr. K Park, Editior. Text Book of Preventive and Social Medicine.
   21st ed. Jabalpur: M/S Banarasidas Bhanot Publication ; 2011. p. 593
- M. Prasad Palthur, S.S. Sajala Palthur, Suresh kumar chitta 2010 Nutraceuticals: A conceptual definition. International journal of pharmacy and pharmaceutical sciences. Vol 4:178. Doi 3

- M. Prasad Palthur, S.S. Sajala Palthur, Suresh kumar chitta 2010 Nutraceuticals: A conceptual definition. International journal of pharmacy and pharmaceutical sciences. Vol 4:178. Doi 3
- Dr. Neeta Mahesh deshpande, Dr. Manasi M deshpande.2017 Kharjur – A review from brihattrayi.World journal of pharmaceutical research. vol 6,doi 9,367-377/2277-7105.
- Kaviraj Ambikadatta shatri, editior.
   Bhaishajjyaratnavali.Chikitsasthan

   Panduroga Chikithsa prakaran.
   Shloka 30. Varanasi: Choukambha Sansrkrit SansthanaPublication; 2001(14<sup>th</sup> edtion). p.378
- 10. CL Ghai, Editior. A Textbook of Practical Physiology. Section 1 Chapter 7.New Delhi: Jaypee brothers medical publishers (P) LTD; 2019.p.34

- 11. Hilbe JM.sigmastat:a second look.Am stat2005;59:187-91.
- 12. Dr. Ghanekar, Commentator. Susrutha Samhita, Sutrasthan. Chapter 14. Delhi (India): Maeherchandlachmandas Publication ; 2010. p. 69
- 13. St, Editior. Anonymous, the Ayurvedic Pharmacopoeia of India (API) Part 4, Vol. IV. New Delhi: Government of India, Ministry of Health and Family Welfare, Dept of I.S.M & H; 2001. P. 63
- 14. Pof.P.V.Sharma, Editior. Dravya Guna Vignana Vol 2. Chapter 14. Varanasi (India): Chaukambha Bharati Acadamey Publication; 1993. p. 759
- 15. Sabnis.M, Editior. Chemistry and Pharmacology of Ayurvedic Medicinal Plants1<sup>st</sup> ed. Varanasi (India): Chaukambha Amarabharati Prakashan; 2006. p. (189.190)

# Table no.1: Nutritional evaluation of dates

Principle	Nutrient value	Percentage of RDA	
		(Recommended Dietary Allowances)	
Energy	277 Kcal	14%	
Carbohydrate	74.97 g	58 %	
Proteins	1.81 g	3 %	
Total fat	0.15 g	< 1 %	
Cholesterol	0 mg	0 %	
Dietary fiber	6.7 gm	18 %	
		Elecrolytes	
Sodium	1 mg	0 %	
Potassium	696 mg	16 %	
		Minerals	

J-RAS

Calcium	64 mg	6.5 %
Copper	0.362 mg	40 %
Iron	0.90 mg	11 %
Magnesium	54 mg	13 %
		Vitamins
Folate	15 µg	4 %
Niacin	1.610 mg	10 %
Pantothenic	0.805 mg	16 %
acid		
Pyridoxine	0.249 mg	19 %
Riboflavin	0.060 mg	4.5 %

Table no.2: Ingredients of Dhatriloha with proportion

Sr no	Name of ingredient	Latin name	Part used	Proportion
<u>1</u>	Amalaki	Embelia ribes Linn	Fruit	Equal
<u>2</u>	Haridra	Curcuma longa Linn	Rhizome	Equal
<u>3</u>	Pippali	Piper longum Linn.	Fruit	Equal
<u>4</u>	Shunthi	Zinziber officinale Linn	Fruit	Equal
<u>5</u>	<u>Maricha</u>	Piper nigrum Linn	Fruit	Equal
<u>6</u>	Lauha Bhasma	incinerated iron	-	Equal

Table no.3: Details of the score adopted for the Rakta Dhatu Kshaya Lakshana

		( \(\\\\)) <	ayurio	9	
Sr no	Symptoms	0	N J <sup>1</sup> R A S	2	3
1	Amla-sheeta Pratharna (craving for sour and cold substances)	Absent	Occasional desire to take either Amla or Sheeta Dravya	Occasional desire to take both <i>Amla</i> and <i>Sheeta Dravya</i>	Always desires for the <i>Amla</i> and <i>Sheeta Dravya</i> that subside on the intake
2	<i>Twaka Parushya</i> (dryness of skin)	No dryness at all –	Dryness in the cold season only	Dryness in other seasons also but only on the exposed parts	Dryness throughout the year but present in the whole body
3	Sira Shaithilya (loss of tone of veins)	Absent	Present occasionally and subside itself after some time	Present occasionally and doesn't subside itself	Present always and subsides on medication

Group	Mean	score	n	0/_	SD	SF	W	T	т	D	Sig
	BT	AT	D	70	30	SL	**	1+	1-	1	Sig
A(n=5)	1.6	0.4	1.2	75	0.447	0.2	15	15	0.00	0.0625	IS
B(n=5)	1.4	0.2	1.2	80	0.447	0.2	15	15	0.00	0.0625	IS

Table no.4: Effect of test drugs on Amla Sheeta Pratharna(craving for sour and cold substances)

A: *Dhatri Loha* treated group, B: *Dhatri loha+Kharjura* treated group, SD: standard deviation, SE: standard error, IS:insignificant

Table no.5: Effect of test drugs on Twaka Parushya (dryness of skin)

Group	Mean	score	п	0/_	SD	SF	W/	T	т	D	Sig
Group	BT	AT	D	/0	30	SE	**	1 +	1-	I	Sig
A(n=5)	1.4	0.6	0.8	60	0.447	0.2	10	10	0	0.125	IS
B(n=5)	1.6	0.4	1.2	80	0.447	0.2	15	15	0	0.0625	IS

A: *Dhatri Loha* treated group, B: *Dhatri Loha+Kharjura* treated group, SD: standard deviation, SE: standard error, IS:insignificant

#### Table no.6: Effect of test drugs on Hb% at 5 days

Crown	<i>B.T.</i>		<i>A.T.</i>		Change		%	(4)
Group	Mean±SEM SD		Mean±SEM SD		Mean±SEM	SD	change	l
A(n=5)	10.900±0.483	1.079	11.140±0.417	0.932	0.240±0.214	0.477	2.39 ↑	-1.124
B(n=5)	10.340 <b>±0.273</b>	0.611	10.760±0.328	0.733	0.420 <b>±0.159</b>	0.356	4.05↑	-2.635

\*\* =p<0.001, \*= p<0.05,  $\downarrow$  = Decrease,  $\uparrow$ = Increase. SEM: standard error of mean Hb%: haemoglobin

percentage, A: Dhatri Loha treated group, B: Dhatri loha+Kharjura treated group,

Table no.7: Effect of test drugs on Hb% 10 at days

Crown	B.T.		А.Т.		Change		%	<b>64</b> 7
Group	Mean±SEM	/lean±SEM SD		Mean±SEM SD		Mean±SEM SD		Υ.
A(n=5)	10.90±0.483	1.079	11.300 <b>±</b> 0.467	1.044	-0.400±0.164	0.367	3.76 ↑	-2.434
B(n=5)	10.340±0.273	0.611	10.980±0.350	0.782	-0.640±0.250	0.559	6.23↑	-2.558

\*\* =p<0.001, \*= p<0.05,↓ = Decrease, ↑= Increase. SEM: standard error of mean Hb%: haemoglobin

percentage, A: Dhatri Loha treated group, B: Dhatri Loha+Kharjura treated group.

Table no.8: Comparison of effect of test drugs on subjective parameters and Hb% by CV test:

Sr no	Parameter	Gr	Group A(n=5)		Gr	oup B(n=	=5)	Better group
		Mean	SD	CV	Mean	SD	CV	
1.	Amla Sheeta	1.2	0.447	0.372	1.2	0.447	0.372	
	Pratharna(craving for sour							Equal
	and cold substances)							
2.	Twaka Parushya (dryness	0.8	0.447	0.558	1.2	0.447	0.372	Group B
	of skin)							
3.	Hb% at 5 days interval	0.24	0.447	1.862	0.42	0.356	0.847	Group B
4.	Hb% at ten days interval	0.4	0.559	1.397	0.64	0.559	0.873	Group B

SD: standard deviation, CV: coefficient of variation, Hb%:percentage of haemoglobin, A: *Dhatri Loha* treated group, B: *Dhatri Loha+Kharjura* treated group.

Group	Mean±SEM	SD	DIFFERNCE	Т	Р	S
A(n=5)	0.240±0.214	0.477	-0.400	-1.216	0.259	IS
B(n=5)	0.640±0.250	0.559				

Table no.9: Comparison of effect on Hb% by unpaired t test

SEM: standard error of mean, SD: standard deviation, IS:insignificant, A: *Dhatri Loha* treated group, B: *Dhatri Loha+Kharjura* treated group

Conflict of Interest: Non

Source of funding: Nil

Cite this article:

Randomised single blind pilot clinical study of kharjura on haemoglobin percentage with

special reference to sushruta's rakta dhatu poshankal."

Supriya P Tonge, Swati Chobhe, Manisha B Walunj

Ayurlog: National Journal of Research in Ayurved Science- 2019; (7) (7): 1 - 11

